

- **Question 1**

1 out of 1 points

In patients with depression, lower doses of SSRIs and SNRIs may be more effective if there is a comorbid anxiety disorder.

Selected Answer:  False

Answers: True  False

- **Question 2**

1 out of 1 points

Match the definitions to one of the following answers: Shave biopsy, punch biopsy, curettage, electrocautery.

Question	Correct Match	Selected Match
Used when knowledge of the depth of the lesion is required	 A. punch biopsy	 A. punch biopsy
Uses a scraping instrument to remove soft and superficial skin lesions	 B. curettage	 B. curettage
Delivery of direct current through a heated metal wire to cause local tissue destruction and hemostasis	 C. electrocautery	 C. electrocautery
Used to remove a lesion when a full-thickness specimen is not required	 D. shave biopsy	 D. shave biopsy

All Answer
Choices

A.

punch biopsy

B.

curettage

C.

electrocautery

D.

shave biopsy

○

- **Question 3**

1 out of 1 points

A manic episode is defined by a period of time (at least 1 week) during which a person's mood is abnormally elevated, expansive, or irritable

Selected



Answer:

True

Answers:



True

Fals

e

- **Question 4**

1 out of 1 points

Palpitations, pounding heart, accelerated heart rate; sweating; trembling or shaking; feeling of choking; chest pain or discomfort; chills or hot flashes; nausea or abdominal distress; feeling dizzy, unsteady, lightheaded, faint; describes patient with what type of mental health diagnosis?

Selected



Answer:

Anxiety