

1. Which of the following are risk factors for neuroleptic malignant syndrome?
Select all that apply

- Age
- Rapid dose escalation
- Parental route of administration
- Higher potency typical antipsychotics

2. Antipsychotic medications provide D2 blockade in the mesocortical pathway causing which of the following effects?

- Reduces negative symptoms
- Increases extra pyramidal symptoms (EPS)
- Causes prolactinemia
- Reduces positive symptoms

3. Phencyclidine (PCP) and Ketamine exert their unique behavioral effects by blocking which of the following receptors?

- GABA type receptors
- Serotonin receptors
- Dopamine receptors
- NMDA type receptors

4. The diagnosis used to describe a syndrome characterized by specific signs and symptoms resulting from recent ingestion or exposure to a substance is known as which of the following?

- Substance intoxication
- Substance withdrawal
- Substance use disorder
- Any of the above

5. Abnormal involuntary movements in a rhythmic pattern affecting face, mouth, tongue, jaw is known as which of the following?

- Akathisia
- Dystonia
- Tardive dyskinesia
- Extrapyramidal side effects

6. A deeply held belief despite evidence to the contrary, lasting at least one month, without prominent hallucinations, with functional impairment that relates to the delusional system

- Schizophrenia
- Delusional disorder
- Brief psychotic disorder
- Schizophreniform disorder

7. Which of the following is true about preoccupation with castration?

- This can be serious and life-threatening if castration is carried out without medical supervision.
- This is consistent with Transvestic Disorder
- This is consistent with other specified paraphilic disorder
- This is consistent with sexual masochism disorder

8. Which of the following is a system that provides a uniform language for describing medical and surgical procedures and diagnostic services rendered by clinicians?

- HIPPA
- DSM-5
- CPT
- DRG

9. A patient on Chlorpromazine states that his orgasm is dry and afterward he will commonly have milky urine. The ARNP realizes which of the following?

- This is likely a harmless response to the Chlorpromazine
- This is abnormal and needs to be evaluated by a specialist
- It is critical to stop the Chlorpromazine immediately
- This has nothing to do with the Chlorpromazine the patient be referred to his PCP for follow-up